arXiv:1304.5698v1 [math-ph] 21 Apr 2013

Liouvillian Propagators, Riccati Equation and Differential Galois Theory

Primitivo B. Acosta-Humánez and Erwin Suazo

To Victoria and Joaquín, propagators of our happiness

Abstract. In this note a Galoisian approach to build propagators through Riccati equations is presented. The main result corresponds to the relationship between the Galois integrability of the linear Schrödinger equation and the virtual solvability of the differential Galois group of its associated characteristic equation. As an application we find the propagator for a generalized harmonic oscillator that has applications describing the process of degenerate parametric amplification in quantum optics and the description the light propagation in a nonlinear anisotropic waveguide.

1. Introduction

The generalized harmonic oscillator has attracted considerable attention over many years in view of their great importance to several advanced quantum problems including Berry's phase, quantization of mechanical of systems and more (see [**11**] and references therein). The fact that in quantum electrodynamics the electromagnetic field can be represented as a set of forced harmonic oscillators makes quadratic Hamiltonians of special interest [**8, 9, 12, 18, 45, 50**]. A method to construct explicit propagators for the linear Schrödinger equation with a time-dependent quadratic Hamiltonian based in solutions of the Riccati equation has been presented in [**7, 8**], and as particular cases the propagators for the free particle, the harmonic oscillator, Caldirola-Kanai and more can be found in a unified manner [**7, 8, 22**].

There has also been an increase interest in the study of Picard-Vessiot Theory, also known as the Galois theory for ordinary linear differential equations, where differential equations of such type are analysed throughout their Galoisian structure. This Galoisian structure depends on the nature of the

---

2010 *Mathematics Subject Classification.* Primary 81Q05; Secondary 12H05.

Key words and phrases. Differential Galois group, Green functions, propagators, Riccati equation.

solutions of the differential equation; for instance, one obtains some kind of solvability (virtual) for the *Galois group* whenever one obtains *Liouvillian solutions*, and in this case one says that the differential equation is *integrable*. This means for example that when one obtains Airy functions, the differential equation is not integrable, while when one obtains Jacobi elliptic functions, the differential equation is integrable, and for instance one gets virtual solvability of its Galois group.

The aim of this paper is to establish a Galoisian approach to the techniques given by Suslov et.al., see [**7, 8, 9, 10**]. To study Liouvillian solutions for linear second order differential equations, as well the integrability of their associated Riccati equations, we use Kovacic algorithm (see [**21**]) and an algebrization procedure (see [**1, 3**]). These tools were applied to study differential equations incoming from physics; in particular the integrability analysis of the one-dimensional linear Schrödinger equation has been studied in [**1, 3**].

As an application of this Galoisian approach we show how to find examples of explicit propagators through Liouvillian solutions for the linear second order differential equation associated to a Riccati equation. As a first example of how to incorporate the Galoisian approach we consider the following (quadratic) generalized harmonic oscillator:

$$\begin{aligned} i\partial_t\psi &= H\psi \qquad (1)\\ H(t) &= \frac{1}{2m}\left(1+\frac{\lambda}{\omega}\cos(2\omega t)\right)p^2+\frac{m\omega^2}{2}\left(1-\frac{\lambda}{\omega}\cos(2\omega t)\right)x^2 \qquad (2)\\ &\quad +\frac{\lambda}{2}\sin(2\omega t)(px+xp), \end{aligned}$$

The oscillator (1)-(2) might be introduced for the first time by Takahashi [**47**] in order to describe the process of degenerate parametric amplification in quantum optics (see also [**23, 24, 26, 27, 35, 41, 47**]). The Hamiltonian (2) had also been considered by Angelow and Trifonov [**4, 5**] in order to describe the light propagation in a nonlinear anisotropic waveguide. In [**11**] the authors constructed the propagator for (1)-(2) using non-periodic solutions of Ince's equation (compare with classical results of periodic solutions [**28**]); they have motivated the study of non-periodic solutions of Ince's equation that seems to be missing in mathematical literature. In fact, they present a non-periodic solution that allows us to write the propagator explicitly; the fact of the solution being non-periodic is fundamental. However, the authors don't get into the details of how they found the solution. In this note we present in detail how to find this non-periodic solution by using a combination of kovacic algorithm [**21**] and an algebrization procedure (see [**1, 3**]). We believe this approach can be extended to the study of propagators of other generalized harmonic oscillators, but here we restrict ourselves to (1)-(2).

This note is organized in the following way:

- Section 2 contains a brief description of the basic theory concerning of construction of explicit propagators using the references [**7, 10, 11, 8, 9, 45, 46, 43, 44**], and a short summary of the Picard-Vessiot theory is also presented, which was written according to [**1, 2, 48**].
- Section 3 contains the main results of this note; it corresponds to a Galoisian approach of propagators and is divided in two parts. The first part is devoted to a theoretical Galoisian approach to propagators starting with Riccati and second order differential equations. The result given here relates the Galois integrability of the linear Schrödinger equation with the virtual solvability of the differential Galois group of its associated characteristic equation. In the second part is presented the Galoisian analysis of second order differential equations to do the construction of new propagators, as well their Green functions through characteristic equations and Riccati equations. Also, we compute the differential Galois group associated to each propagator, which corresponds to the differential Galois group of the characteristic equation.
- It has been included an appendix, written according to [**1, 2, 3**], which describes the Kovacic algorithm and the Hamiltonian algebrization, tools used to compute the propagators in Section 3.

For suitability, along this paper ∂_x denotes $\frac{\partial}{\partial x}$, for higher order derivation ∂_x^n denotes $\frac{\partial^n}{\partial x^n}$ and by $\partial_t a(0)$ we mean $\partial_t a(t)|_{t=0}$.

2. Theoretical Background

2.1. Differential Galois theory. Theoretical aspects of Galoisian approach to differential equations are summarised in this subsection which is based on the references [**19, 20, 21, 30, 48**]. Moreover, applications of differential Galois theory to mathematical physics can be found in [**1, 2, 3, 32, 33, 42**]. The Galois theory of differential equations, also called Differential Galois Theory has been developed by Picard, Vessiot, Kolchin and currently by a lot of researchers. In particular, we focus in the Galois theory of linear differential equations, also known as Picard-Vessiot theory. Following [**3**] and also [**1**], we present here an algebraic model for functions and the corresponding Galois theory.

Definition 1 (Differential Fields). . *Let K be a commutative field of characteristic zero. A* derivation *of K is a map $\partial_x : K \to K$ satisfying $\partial_x(a+b) = \partial_x a + \partial_x b$ and $\partial_x(ab) = \partial_x a \cdot b + a \cdot \partial_x b$ for all $a, b \in K$. We then say that (K, ∂_x) (or just K, when there is no ambiguity) is a differential field with the derivation ∂_x.*

We assume that K contains an element x such that $\partial_x(x) = 1$. Let $\mathcal{C}$ denote the field of constants of K:

$$\mathcal{C} = \{c \in K | \partial_x c = 0\}.$$

It is also of characteristic zero and will be assumed to be algebraically closed. The *coefficient field* for a differential equation is defined as the smallest differential field containing all the coefficients of the equation.
Due to we will mostly analyze second order linear homogeneous differential equations, i.e equations of the form

$$\mathcal{L} := \partial_x^2 y + a\partial_x y + by = 0, \quad a, b \in K.$$

so the rest of the theory will be explained in this context.

Definition 2 (Picard-Vessiot Extension). *Let L be a differential field containing K (a differential extension of K). We say that L is a* Picard-Vessiot *extension of K for $\mathcal{L}$ if there exist two linearly independent $y_1, y_2 \in L$ solutions of $\mathcal{L}$ such that $L = K\langle y_1, y_2\rangle$ (i.e $L = K(y_1, y_2, \partial_x y_1, \partial_x y_2)$) and L and K has the same field of constants $\mathcal{C}$.*

In what follows, we choose a Picard-Vessiot extension and the term "solution of $\mathcal{L}$" will mean "solution of $\mathcal{L}$ in L". So any solution of $\mathcal{L}$ is a linear combination (over $\mathcal{C}$) of y_1 and y_2.

A K-automorphism σ of the Picard-Vessiot extension L is called a differential automorphism if it leaves K fixed and commutes with the derivation. This means that $\sigma(\partial_x a) = \partial_x(\sigma(a))$ for all $a \in L$ and $\forall a \in K$, $\sigma(a) = a$.

Definition 3 (Differential Galois Groups). *The group of all differential automorphisms of L over K is called the differential Galois group of L over K and is denoted by* $\mathrm{DGal}(L/K)$.

Given $\sigma \in \mathrm{DGal}(L/K)$, we see that $\{\sigma y_1, \sigma y_2\}$ are also solutions of $\mathcal{L}$. Hence there exists a matrix

$$A_\sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}(2, \mathbb{C}),$$

such that

$$\sigma(\begin{pmatrix} y_1 & y_2 \end{pmatrix}) = \begin{pmatrix} \sigma(y_1) & \sigma(y_2) \end{pmatrix} = \begin{pmatrix} y_1 & y_2 \end{pmatrix} A_\sigma.$$

As σ commutes with the derivation, this extends naturally to an action on a fundamental solution matrix of the companion first order system associated with $\mathcal{L}$.

$$\sigma\left(\begin{pmatrix} y_1 & y_2 \\ \partial_x y_1 & \partial_x y_2 \end{pmatrix}\right) = \begin{pmatrix} \sigma(y_1) & \sigma(y_2) \\ \sigma(\partial_x y_1) & \sigma(\partial_x y_2) \end{pmatrix} = \begin{pmatrix} y_1 & y_2 \\ \partial_x y_1 & \partial_x y_2 \end{pmatrix} A_\sigma.$$

This defines a faithful representation $\mathrm{DGal}(L/K) \to \mathrm{GL}(2, \mathbb{C})$ and it is possible to consider $\mathrm{DGal}(L/K)$ as a subgroup of $\mathrm{GL}(2, \mathbb{C})$. It depends on the choice of the fundamental system $\{y_1, y_2\}$, but only up to conjugacy.
Recall that an algebraic group G is an algebraic manifold endowed with a group structure. Let $\mathrm{GL}(n, \mathbb{C})$ denote, as usual, the set of invertible $n \times n$ matrices with entries in $\mathbb{C}$ (and $\mathrm{SL}(n, \mathbb{C})$ be the set of matrices with determinant equal to 1). A linear algebraic group will be a subgroup of $\mathrm{GL}(n, \mathbb{C})$ equipped with a structure of algebraic group. One of the fundamental results of the Picard-Vessiot theory is the following theorem.

Theorem 4. [**20**] *The differential Galois group* $\mathrm{DGal}(L/K)$ *is an algebraic subgroup of* $\mathrm{GL}(2,\mathbb{C})$.

In fact, the differential Galois group measures the algebraic relations between the solutions (and their derivatives). It is sometimes viewed as the object which should tell "what algebra sees of the dynamics of the solutions".

In an algebraic group G, the largest connected algebraic subgroup of G containing the identity, noted G°, is a normal subgroup of finite index. It is often called the connected component of the identity. If $G = G^0$ then G is a *connected group*. Furthermore, when G^0 satisfies some property, we say that G virtually satisfies this property. For example, virtually solvability of G means solvability of G^0.

Theorem 5 (Lie-Kolchin Theorem). *Let* $G \subseteq \mathrm{GL}(2,\mathbb{C})$ *be a virtually solvable group. Then* G^0 *is triangularizable, i.e it is conjugate to a subgroup of upper triangular matrices.*

The following corresponds to the list of the *Algebraic Subgroups of* $\mathrm{SL}(2,\mathbb{C})$, see [**20, 21, 48**] for further details.

- *Reducible subgroups*. These are the groups which leave a non-trivial subspace of V invariant. They are classified in two categories.
    - *Diagonal groups:*
      the identity group:
      $$\{e\} = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\},$$
      the $n-$roots group:
      $$\mathbb{G}^{[n]} = \left\{ \begin{pmatrix} c & 0 \\ 0 & c^{-1} \end{pmatrix}, \quad c^n = 1 \right\},$$
      the multiplicative group:
      $$\mathbb{G}_m = \left\{ \begin{pmatrix} c & 0 \\ 0 & c^{-1} \end{pmatrix}, \quad c \in \mathbb{C}^* \right\}.$$
    - *Triangular groups:*
      the additive group:
      $$\mathbb{G}_a = \left\{ \begin{pmatrix} 1 & d \\ 0 & 1 \end{pmatrix}, \quad d \in \mathbb{C} \right\},$$
      the $n-$quasi-roots:
      $$\mathbb{G}^{\{n\}} = \left\{ \begin{pmatrix} c & d \\ 0 & c^{-1} \end{pmatrix}, \quad c^n = 1, \quad d \in \mathbb{C} \right\},$$
      the Borel group:
      $$\mathbb{B} = \left\{ \begin{pmatrix} c & d \\ 0 & c^{-1} \end{pmatrix}, \quad c \in \mathbb{C}^*, \quad d \in \mathbb{C} \right\}.$$
- *Irreducible subgroups*

- The infinite dihedral group (also called meta-abelian group):

$$\mathbb{D}_\infty = \left\{ \begin{pmatrix} c & 0 \\ 0 & c^{-1} \end{pmatrix}, \quad c \in \mathbb{C}^* \right\} \cup \left\{ \begin{pmatrix} 0 & d \\ -d^{-1} & 0 \end{pmatrix}, \quad d \in \mathbb{C}^* \right\}$$

  and its finite subgroups $\mathbb{D}_{2n}$ (where c and d spans the n-th roots of unity).
- Finite irreducible (primitive) groups:
  the tetrahedral group $A_4^{\mathrm{SL}_2}$ of order 24,
  the octahedral group $S_4^{\mathrm{SL}_2}$ of order 48,
  and the icosahedral group $A_5^{\mathrm{SL}_2}$ of order 120.

Definition 6 (Integrability of $\mathcal{L}$). *We say that the linear differential equation $\mathcal{L}$ is (Liouville) integrable if the Picard-Vessiot extension $L \supset K$ is obtained as a tower of differential fields $K = L_0 \subset L_1 \subset \cdots \subset L_m = L$ such that $L_i = L_{i-1}(\eta)$ for $i = 1, \ldots, m$, where either*

(1) *η is* algebraic *over L_{i-1}, that is η satisfies a polynomial equation with coefficients in L_{i-1}.*
(2) *η is* primitive *over L_{i-1}, that is $\partial_x \eta \in L_{i-1}$.*
(3) *η is* exponential *over L_{i-1}, that is $\partial_x \eta/\eta \in L_{i-1}$.*

We remark that the usual terminology in differential algebra for integrable equations is that the corresponding Picard-Vessiot extensions are called *Liouvillian*. The following theorem is due to Kolchin.

Theorem 7. *The equation $\mathcal{L}$ is integrable if and only if* $\mathrm{DGal}(L/K)$ *is virtually solvable.*

The following result is taken from [**2**] and corresponds to the transformations involving Riccati equations and second order differential equations.

Proposition 8 (Riccati transformations, [**2**]). *Let K be a differential field and let consider functions $a_0(x)$, $a_1(x)$, $a_2(x)$, $r(x)$, $\rho(x)$, $b_0(x)$, $b_1(x)$ belonging to K that, for simplicity, will be denoted without their explicit dependence on x. Consider now the following forms associated to any second order differential equation (ode) and Riccati equation:*

(*i*) *Second order ode (in general form):*

$$\partial_x^2 y + b_1 \partial_x y + b_0 y = 0. \tag{3}$$

(*ii*) *Second order ode (in reduced form):*

$$\partial_x^2 \xi = \rho \xi. \tag{4}$$

(*iii*) *Riccati equation (in general form):*

$$\partial_x v = a_0 + a_1 v + a_2 v^2, \quad a_2 \neq 0. \tag{5}$$

(*iv*) *Riccati equation (in reduced form):*

$$\partial_x w = r - w^2, \tag{6}$$

Then, there exist transformations T, B, S and R leading some of these equations into the other ones, as showed in the following diagram:

$$\begin{tikzcd}
\partial_x v = a_0 + a_1 v + a_2 v^2 \arrow[r, "T"] \arrow[d, "B"'] & \partial_x w = r - w^2 \arrow[d, "R"] \\
\partial_x^2 y + b_1 \partial_x y + b_0 y = 0 \arrow[r, "S"] & \partial_x^2 \xi = \rho \xi.
\end{tikzcd}$$

The new independent variables are defined by means of

$$T:\ v = -\left(\frac{\partial_x a_2}{2a_2^2} + \frac{a_1}{2a_2}\right) - \frac{1}{a_2} w, \qquad B:\ v = -\frac{1}{a_2}\frac{\partial_x y}{y},$$

$$S:\ y = \xi e^{-\frac{1}{2}\int b_1 dx}, \qquad R:\ w = \frac{\partial_x \xi}{\xi},$$

and the functions r, ρ, b_0 and b_1 are given by

$$r = \frac{1}{\beta}\left(a_0 + a_1\alpha + a_2\alpha^2 - \partial_x\alpha\right), \tag{7}$$

$$\alpha = -\left(\frac{\partial_x a_2}{2a_2^2} + \frac{a_1}{2a_2}\right), \quad \beta = -\frac{1}{a_2}, \tag{8}$$

$$b_1 = -\left(a_1 + \frac{\partial_x a_2}{a_2}\right), \quad b_0 = a_0 a_2, \tag{9}$$

$$\rho = r = \frac{b_1^2}{4} + \frac{\partial_x b_1}{2} - b_0. \tag{10}$$

Remark 9. *From Proposition 8, it is recovered the well known result in differential Galois theory (see for example* [**48**]*): Riccati equation has an algebraic solution over the differential field K if and only if its associated second order differential equation has two independent Liouvillian solutions (the differential Galois group of the second order differential equation is virtually solvable).*

2.2. Propagators and Green Functions. In this section, as well in the rest of the paper, we follow [**10**] considering the one-dimensional time-dependent Schrödinger equation for an harmonic oscillator

$$i\partial_t\psi = H\psi, \quad H = a(t)p^2 + b(t)x^2 + c(t)(px + xp), \quad p = i\partial_x. \tag{11}$$

The Schrödinger equation (11) can be written as

$$i\partial_t\psi = -a(t)\,\partial_x^2\psi + b(t)\,x^2\psi - ic(t)\,\psi - i\frac{c(t)}{2}x\partial_x\psi. \tag{12}$$

We start considering the Riccati equation

$$\partial_t\alpha + b(t) + 2c(t)\,\alpha + 4a(t)\,\alpha^2 = 0, \tag{13}$$

where $a(t)$, $b(t)$ and $c(t)$ are elements of a differential field K, with coefficient field $\mathbb{C}$. By Proposition 8 we can transform the Riccati equation (13) through the change of variable

$$\alpha (t) = \frac{1}{4a (t)} \frac{\partial_t \mu (t)}{\mu (t)} - \frac{c(t)}{2a(t)}, \tag{14}$$

into the second order differential equation

$$\partial_t^2 \mu - \tau(t) \partial_t \mu + 4\sigma(t) \mu = 0. \tag{15}$$

Moreover, by Remark 9, the differential Galois group of the differential equation (15) is virtually solvable if and only if the Riccati equation (13) has an algebraic solution over the differential field K. Furthermore, by Proposition 8, departing from the differential equation (15) we can arrive at the Riccati equation (13) through changes of variables. The following lemmas show how can we construct propagators based on explicit solutions in (13) and (15).

Lemma 10. [**7, 10, 44**] *Given $a(t), b(t)$ and $c(t)$ piecewise continuous, there exists an interval I of time where the following (Riccati-type) system*

$$\partial_t \alpha + b (t) + 2c (t) \alpha + 4a (t) \alpha^2 = 0, \tag{16}$$

$$\partial_t \beta + (c (t) + 4a (t) \alpha (t)) \beta = 0, \tag{17}$$

$$\partial_t \gamma + a (t) \beta^2 (t) = 0, \tag{18}$$

has as a fundamental solution in terms of solutions of the following (characteristic) equation

$$\partial_t^2 \mu - \tau (t) \partial_t \mu + 4\sigma (t) \mu = 0 \tag{19}$$

$$\tau (t) = \frac{\partial_t a}{a} - 4c, \qquad \sigma (t) = ab + \frac{c^2}{2} + \frac{c}{4} \left(\frac{\partial_t a}{a} - \frac{\partial_t c}{c} \right) \tag{20}$$

given by:

$$\alpha_0 (t) = \frac{1}{4a (t)} \frac{\partial_t \mu_0 (t)}{\mu_0 (t)} - \frac{c (t)}{4a (t)}, \tag{21}$$

$$\beta_0 (t) = - \frac{1}{\mu_0 (t)}, \tag{22}$$

$$\gamma_0 (t) = \frac{1}{2\mu_1 (0)} \frac{\mu_1 (t)}{\mu_0 (t)} + \frac{d (0)}{2a (0)} \tag{23}$$

provided that μ_0 and μ_1 are standard solutions of (19)-(20) with $\mu_0 (0) = 0$, $\partial_t \mu_0 (0) = 2a (0) \neq 0$, and $\mu_1 (0) \neq 0$, $\partial_t \mu_1 (0) = 0$, and $\delta_0 (0) = -\varepsilon_0 (0) = g (0) / (2a (0))$, $\kappa_0 (0) = 0$. Further, the following asymptotics hold:

$$\alpha_0 (t) = \frac{1}{4a (0) t} - \frac{c (0)}{4a (0)} - \frac{\partial_t a(0)}{8a^2(0)} + O(t), \tag{24}$$

$$\beta_0 (t) = - \frac{1}{2a (0) t} + \frac{\partial_t a(0)}{4a^2(0)} + O(t), \tag{25}$$

$$\gamma_0 (t) = \frac{1}{4a (0) t} + \frac{c (0)}{4a (0)} - \frac{\partial_t a(0)}{8a^2(0)} + O(t) \tag{26}$$

as $t \to 0$, with $a(t), b(t)$ and $c(t)$ sufficiently smooth.

Lemma 11. [**7, 10, 44, 45**] *The Green function, or Feynman's propagator, corresponding to the Schrödinger equation 11 can be obtained as*

$$\psi = G\left(x,y,t\right) = \frac{1}{\sqrt{2\pi i \mu_0\left(t\right)}}\ e^{i\left(\alpha_0(t)x^2+\beta_0(t)xy+\gamma_0(t)y^2\right)}, \tag{27}$$

where $\alpha_0\left(t\right)$, $\beta_0\left(t\right)$ *and* $\gamma_0\left(t\right)$ *are solutions of the Riccati-type system. Then the superposition principle allows us to solve the corresponding Cauchy initial value problem:*

$$\psi(x,t) = \int_{-\infty}^{\infty} G(x,y,t)\psi(y,0)dy$$

for suitable data $\psi(x,0)= \varphi(x)$. *Further as* $t \to 0$,

$$\begin{aligned} G\left(x,y,t\right) \quad \sim \quad & \frac{1}{\sqrt{2\pi i a(0)t}} \exp\left[i\frac{\left(x-y\right)^2}{4a(0)t}\right] \\ & \times \exp\left[-i\left(\frac{\partial_t a(0)}{8a^2(0)}\left(x-y\right)^2 + \frac{c\left(0\right)}{4a\left(0\right)}\left(x-y\right)^2\right)\right]. \end{aligned}$$

3. Galoisian approach to Propagators

In this section we apply the Picard-Vessiot theory in the context of propagators.

We are interested in *Liouvillian solutions* of the linear Schrödinger equation (11); that is, the so-called *Liouvillian propagator* (27) is obtained through Liouvillian functions. In this way, we can give a Galoisian formulation for this kind of integrability.

Definition 12. *Linear Schrödinger equation (11) is integrable in Galoisian sense (Galois integrable), when it has a Liouvillian propagator (27).*

Theorem 13 (Galoisian approach to LSE). *Linear Schrödinger equation (11) is Galois integrable if and only if the differential Galois group of the characteristic equation (15) is virtually solvable.*

Proof. Consider K as the differential field of the characteristic equation. Let us suppose (see Lemma 10) that μ_0 is a Liouvillian solution of the characteristic equation; then μ_1 is Liouvillian too. This implies that the differential Galois group of the characteristic equation is virtually solvable and α is an algebraic solution of the Riccati equation over K. Now, using μ_0 and μ_1 in β and γ such as in Lemma 10, we see that they are Liouvillian functions over K and therefore the propagator is Liouvillian. Thus, by Definition 12 we get that the Schrödinger equation is Galois integrable. Conversely, assuming that the Schrödinger equation is Galois integrable we see that it has a Liouvillian propagator that can be obtained through an algebraic solution of the Riccati equation over K. For instance, there exists a Liouvillian solution μ_0 of the characteristic equation which implies that its differential Galois group is virtually solvable. $\square$

Remark 14. *In virtue of Theorem 13 we can construct many Liouvillian propagators through integrable second order differential equations over a differential field as well by algebraic solutions, over such differential field, of Riccati equations. This is the practical aim of this paper, which will be given in Section 3.1.*

3.1. New Propagators From Differential Galois Theory.

In this section we use the Galoisian approach to LSE, Theorem 13, to generalize examples introduced in [**22**] and find the explicit solution presented by Cordero-Suslov in [**10**], as well as to introduce new propagators. The starting point is the knowledge of the integrability, in the Picard-Vessiot sense, of the characteristic equation, or equivalently the existence of an algebraic solution over its differential field of its associated Riccati equation. To obtain the solutions of the characteristic equations we apply two practical tools of differential Galois theory: *Kovacic Algorithm* and *Hamiltonian Algebrization*, see Appendix A.

3.1.1. *Characteristic equation* $\partial_t^2\mu + t^n\mu = 0$. We consider in this equation as differential field to $K = \mathbb{C}(t)$. This is a generalization of the case $n = 1$, which was presented in [**22**]. However, the case $n = 1$ does not correspond to Galoisian integrability of the characteristic equation: the solutions are not Liouvillian due to the fact that they are Airy functions. Now, we study the integrability, in Galoisian sense, of this equation through Kovacic Algorithm. As in [**42**], we obtain three conditions for n to get virtual solvability of the differential Galois group. It follows the construction of the propagators related with these integrability conditions.

(1) Let's consider the characteristic equation $\partial_t^2\mu + \mu = 0$ ($n = 0$). A basis of solutions is given by $\mathcal{B} = \{\sin t, \cos t\}$. Thus, the Picard-Vessiot extension is given by $L = \mathbb{C}(t, e^{it})$ and for instance the differential Galois group is $\mathrm{DGal}(L/K) = \mathbb{G}_m$. By Proposition 8, through the change of variable $\alpha = \partial_t\mu/\mu$, we obtain the Riccati equation (13), where $a = 1/4$, $b = 1$ and $c = 0$. Considering $\mu_0(t) = \lambda_1 \sin t$ and $\mu_1(t) = \lambda_2 \cos t$, we see that the conditions of Lemma 10 are satisfied when $\lambda_1 = 1/2$. Furthermore, λ_2 must be 2 to get $W(\mu_0, \mu_1) = 1$. In this way, by Theorem 13 the Schrödinger equation is Galois integrable and its propagator is Liouvillian.

(2) Let's consider the characteristic equation $\partial_t^2\mu + \mu/t^2 = 0$ ($n = -2$). A basis of solutions is given by $\mathcal{B} = \{t^{m+1}, t^{-m}\}$, being $m = (-1 \pm \sqrt{5})/2$. Since $m \notin \mathbb{Q}$, the Picard-Vessiot extension is $L = \mathbb{C}(t, t^m)$ and therefore $\mathrm{DGal}(L/K) = \mathbb{G}_m$. In [**1**] there is a complete study of the Galoisian structure of this equation.

(3) The characteristic equation $\partial_t^2\mu + \mu/t^4 = 0$ ($n = -4$). A basis of solutions is given by $\mathcal{B} = \{t\cos(1/t), t\sin(1/t)\}$. Thus, the Picard-Vessiot extension is $L = \mathbb{C}(t, e^{i/t})$ and therefore $\mathrm{DGal}(L/K) = \mathbb{G}_m$. Here, initial conditions for the characteristic equation are satisfied when $t \to 0^+$, that is, $\mu_0(t) \to 0$ when $t \to 0^+$.

3.1.2. *Characteristic equation* $\partial_t^2\mu + 2\lambda\tan(\lambda t)\partial_t\mu - 2\mu = 0$. This equation is a particular case of the Ince's equation

$$\partial_t^2\mu + \frac{2\lambda\,\omega\,\sin(2\,\omega\,t)}{\omega + \lambda\,\cos(2\,\omega\,t)}\partial_t\mu + \frac{\omega^3 - 3\,\omega\,\lambda^2 - \left(\omega^2\lambda + \lambda^3\right)\cos(2\,\omega\,t)}{\omega + \lambda\,\cos(2\,\omega\,t)}\mu = 0$$

when $\lambda = \omega$. The general case corresponds to a parametric oscillator introduced in [**11**], where the case $\lambda = \omega = 1$ was completely analyzed. To avoid triviality, we consider $\lambda \neq 0$ for the characteristic equation which will be analysed with differential Galois theory. To do this, we apply the hamiltonian algebrization process and Kovacic's algorithm on the characteristic equation, and so we obtain the solutions as well its differential Galois group. Afterward, we construct the corresponding propagator associated to this characteristic equation. We consider $K = \mathbb{C}(\tan\lambda t)$ as the differential field of the characteristic equation.

We first consider the hamiltonian change of variable $\tau = \tan\lambda t$ obtaining

$$\partial_\tau^2\hat{\mu}(\tau) + \frac{4\tau}{1+\tau^2}\partial_\tau\hat{\mu}(\tau) - \frac{2}{(1+\tau^2)^2}\hat{\mu}(\tau) = 0, \tag{28}$$

i.e., the algebraic form of the characteristic equation, being $\widehat{K} = \mathbb{C}(\tau)$ its differential field which is isomorphic to $K = \mathbb{C}(\tan\lambda t)$. In order to apply the Kovacic algorithm we should reduce the equation (28) through the change of dependent variable

$$\hat{\mu}(\tau) = y\exp\left(-\frac{1}{2}\int_0^\tau \frac{4s}{1+s^2}ds\right) = \frac{y}{1+\tau^2}.$$

Thus we obtain the reduced form

$$\partial_\tau^2 y = ry, \quad r = \frac{2\tau^2+4}{(1+\tau^2)^2} = \frac{2}{1+\tau^2} + \frac{2}{(1+\tau^2)^2}, \tag{29}$$

where the Picard-Vessiot extension is the same as in (28), denoted by $\widehat{L}$, and for instance the differential Galois group will be the same for both equations due to the fact that their coefficients are rational functions in τ, i.e., $\widehat{K} = \mathbb{C}(\tau)$.

Applying Kovacic's algorithm we see that $\Gamma = \{i, -i, \infty\}$, $\circ r_i = \circ r_{-i} = \circ r_\infty = 2$, which implies that equation (29) can fall in cases 1, 2, 3 or 4 of the algorithm. We begin analyzing case one.

Case 1. According to step 1, we check conditions (c_2) and (∞_2) to obtain $b_i = b_{-i} = -\frac{1}{2}$ and $b_\infty = 2$. In this way, $\alpha_i^\pm = \alpha_{-i}^\pm = \frac{1\pm i}{2}$, $\alpha_\infty^\pm = \frac{1\pm 3}{2}$,

$\sqrt{r}_i = \sqrt{r}_{-i} = \sqrt{r}_\infty = 0$. By step 2 we obtain $D = \{1\}$ through two different options: $1 = \alpha_\infty^+ - \alpha_i^+ - \alpha_{-i}^-$ and $1 = \alpha_\infty^+ - \alpha_i^- - \alpha_{-i}^+$; for instance we have two possibilities for ω: $\omega_1 = \frac{1}{\tau-i} - \frac{1}{\tau+i}$ and $\omega_2 = \frac{1}{\tau+i} - \frac{1}{\tau-i}$, that is, $\omega_1^2 = \omega_2^2$. Following step 3 we see that there does not exist a polynomial $P_1(\tau) = \tau + a_0$ satisfying the algebraic equation (32). In this way, we should go to the case 2 of Kovacic algorithm due to $\mathrm{DGal}(\widehat{L}/\widehat{K})$ not being a subgroup of the Borel group.

Case 2. According to step 1, we check conditions (c_2) and (∞_2) to obtain $b_i = b_{-i} = -\frac{1}{2}$ and $b_\infty = 2$. In this way, $E_i = E_{-i} = \{2\}$, $E_\infty = \{-4, 2, 8\}$. By step 2 we obtain $D = \{2\}$ only for $e_\infty = 8$. In this way, we obtain $\theta = \frac{1}{\tau-i} + \frac{1}{\tau+i}$. Following step 3 we obtain the polynomial $P_2(\tau) = \tau^2 - 1$ satisfying the algebraic equation (33), which leads us to obtain $\phi = \frac{1}{\tau-i} + \frac{1}{\tau+i} + \frac{2\tau}{\tau^2-1}$. In this way, solving the algebraic equation

$$\omega^2 + \phi\omega + \frac{1}{2}(\partial_\tau \phi + \phi^2 - 2r) = 0,$$

we obtain two solutions for ω:

$$\omega_- = 2\,\frac{\tau^2 - \tau + 1}{\tau^3 - \tau^2 + \tau - 1}, \quad \omega_+ = 2\,\frac{\tau^2 + \tau + 1}{\tau^3 + \tau^2 + \tau + 1}.$$

In this way, due to $y = e^{\int \omega d\tau}$, we have the general solution of (29):

$$y = C_1 e^{-\arctan\tau}(\tau - 1)\sqrt{1+\tau^2} + C_2 e^{\arctan\tau}(\tau + 1)\sqrt{1+\tau^2},$$

for instance $\mathrm{DGal}(\widehat{L}/\widehat{K}) = \mathbb{D}_\infty$. Now, the general solution for equation (28) is given by

$$\widehat{\mu}(\tau) = C_1 \frac{e^{-\arctan\tau}(\tau - 1)}{\sqrt{1+\tau^2}} + C_2 \frac{e^{\arctan\tau}(\tau + 1)}{\sqrt{1+\tau^2}};$$

for instance the differential Galois group for the algebrized characteristic equation (28) is also the dihedral infinite group $\mathbb{D}_\infty$. Recalling that $\tau = \tan\lambda t$, we get the general solution of the characteristic equation

$$\mu(t) = C_1 e^{-\lambda t}(\sin\lambda t - \cos\lambda t) + C_2 e^{\lambda t}(\sin\lambda t + \cos\lambda t),$$

which can also be written as

$$\mu(t) = (C_1 + C_2)(\sinh\lambda t\cos\lambda t + \cosh\lambda t + \sin\lambda t$$
$$+(C_2 - C_1)\sinh\lambda t\sin\lambda t + \cosh\lambda t\cos\lambda t,$$

and its differential Galois group is also the dihedral infinite group, i.e.,$\mathrm{DGal}(L/K) = \mathbb{D}_\infty$. Now, we find $\mu_0(t)$ and $\mu_1(t)$ satisfying the conditions of Lemma 10

$$\mu_0(t) = \sinh\lambda t\cos\lambda + \cosh\lambda t\sin\lambda t, \quad C_1 = C_2 = \frac{1}{2},$$

$$\mu_1(t) = \sinh\lambda t\cos\lambda + \cosh\lambda t\sin\lambda t, \quad -C_1 = C_2 = \frac{1}{2},$$

which for $\lambda = 1$ corresponds to the solutions given in [**11**]. Finally, the propagator of the Schrödinger equation for this characteristic equation are given by

$$\begin{aligned} G\left(x,y,t\right) &= \frac{1}{\sqrt{2\pi i(\cos\lambda t\sinh\lambda t+\sin\lambda t\cosh\lambda t)}} \\ &\quad\times\exp\left[\frac{(x^2-y^2)\sin\lambda t\sinh\lambda t+2xy-(x^2-y^2)\cos\lambda t\cosh\lambda t}{2i(\cos\lambda t\sinh\lambda t+\sin\lambda t\cosh\lambda t)}\right] \end{aligned} \tag{30}$$

Final Remarks

This paper is an starting point to study the integrability of partial differential equations in a more general sense through differential Galois theory. With this approach we studied the linear Schrödinger equation corresponding to a generalized (quadratic) harmonic oscillator obtaining similar results to [**11, 22**].

Acknowledgments

The first author is partially supported by the MICIIN/FEDER grant number MTM2009–06973, by the Generalitat de Catalunya grant number 2009SGR859 and by DIDI – Universidad del Norte. The second author is granted by Banco Santander. The authors thank to Carles Simó and Juan J. Morales-Ruiz by their useful comments and suggestions.

Appendix A. Kovacic Algorithm and Hamiltonian Algebrization

This appendix contains the basic information related with two practical tools of differential Galois theory, which have been used to compute the propagators in Section 3.

A.1. Kovacic algorithm. Kovacic in 1986 (see [**21**]) introduced an algorithm to solve the differential equation $\partial_x^2\zeta = r\zeta$, where $r \in \mathbb{C}(x)$.

Each case in Kovacic's algorithm is related with each one of the algebraic subgroups of $\mathrm{SL}(2,\mathbb{C})$ and the associated Riccati equation

$$\partial_x v = r - v^2 = \left(\sqrt{r}-v\right)\left(\sqrt{r}+v\right), \quad v = \frac{\partial_x\zeta}{\zeta}.$$

There are four cases in Kovacic's algorithm. Only for cases 1, 2 and 3 we can solve the differential equation, but for the case 4 the differential equation is not integrable. It is possible that Kovacic's algorithm can provide us only one solution (ζ_1), so that we can obtain the second solution (ζ_2) through

$$\zeta_2 = \zeta_1\int\frac{dx}{\zeta_1^2}. \tag{31}$$

For the differential equation given by

$$\partial_x^2\zeta = r\zeta, \qquad r = \frac{s}{t}, \quad s,t \in \mathbb{C}[x],$$

we use the following notations.

1. Denote by Γ' be the set of (finite) poles of r, $\Gamma' = \{c \in \mathbb{C} : t(c) = 0\}$.
2. Denote by $\Gamma = \Gamma' \cup \{\infty\}$.
3. By the order of r at $c \in \Gamma'$, $\circ(r_c)$, we mean the multiplicity of c as a pole of r.
4. By the order of r at ∞, $\circ(r_\infty)$, we mean the order of ∞ as a zero of r. That is $\circ(r_\infty) = \deg(t) - \deg(s)$.

Case 1. In this case $[\sqrt{r}]_c$ and $[\sqrt{r}]_\infty$ means the Laurent series of $\sqrt{r}$ at c and the Laurent series of $\sqrt{r}$ at ∞ respectively. Furthermore, we define $\varepsilon(p)$ as follows: if $p \in \Gamma$, then $\varepsilon(p) \in \{+,-\}$. Finally, the complex numbers $\alpha_c^+, \alpha_c^-, \alpha_\infty^+, \alpha_\infty^-$ will be defined in the first step. If the differential equation has no poles it only can fall in this case.

Step 1. Search for each $c \in \Gamma'$ and for ∞ the corresponding situation as follows:

(c_0): If $\circ(r_c) = 0$, then

$$\left[\sqrt{r}\right]_c = 0, \quad \alpha_c^\pm = 0.$$

(c_1): If $\circ(r_c) = 1$, then

$$\left[\sqrt{r}\right]_c = 0, \quad \alpha_c^\pm = 1.$$

(c_2): If $\circ(r_c) = 2$, and

$$r = \cdots + b(x-c)^{-2} + \cdots, \quad \text{then}$$

$$\left[\sqrt{r}\right]_c = 0, \quad \alpha_c^\pm = \frac{1 \pm \sqrt{1+4b}}{2}.$$

(c_3): If $\circ(r_c) = 2v \geq 4$, and

$$r = (a\,(x-c)^{-v} + ... + d\,(x-c)^{-2})^2 + b(x-c)^{-(v+1)} + \cdots, \quad \text{then}$$

$$\left[\sqrt{r}\right]_c = a\,(x-c)^{-v} + ... + d\,(x-c)^{-2}, \quad \alpha_c^\pm = \frac{1}{2}\left(\pm\frac{b}{a} + v\right).$$

(∞_1): If $\circ(r_\infty) > 2$, then

$$\left[\sqrt{r}\right]_\infty = 0, \quad \alpha_\infty^+ = 0, \quad \alpha_\infty^- = 1.$$

(∞_2): If $\circ(r_\infty) = 2$, and $r = \cdots + bx^2 + \cdots$, then

$$\left[\sqrt{r}\right]_\infty = 0, \quad \alpha_\infty^\pm = \frac{1 \pm \sqrt{1+4b}}{2}.$$

(∞_3): If $\circ(r_\infty) = -2v \leq 0$, and

$$r = (ax^v + ... + d)^2 + bx^{v-1} + \cdots, \quad \text{then}$$

$$\left[\sqrt{r}\right]_\infty = ax^v + ... + d, \quad \textit{and} \quad \alpha_\infty^\pm = \frac{1}{2}\left(\pm\frac{b}{a} - v\right).$$

Step 2. Find $D \neq \emptyset$ defined by

$$D = \left\{ n \in \mathbb{Z}_+ : n = \alpha_\infty^{\varepsilon(\infty)} - \sum_{c\in\Gamma'} \alpha_c^{\varepsilon(c)}, \forall \left(\varepsilon\left(p\right)\right)_{p\in\Gamma} \right\}.$$

If $D = \emptyset$, then we should start with the case 2. Now, if $\mathrm{Card}(D) > 0$, then for each $n \in D$ we search $\omega \in \mathbb{C}(x)$ such that

$$\omega = \varepsilon\left(\infty\right)\left[\sqrt{r}\right]_\infty + \sum_{c\in\Gamma'} \left(\varepsilon\left(c\right)\left[\sqrt{r}\right]_c + \alpha_c^{\varepsilon(c)}(x-c)^{-1} \right).$$

Step 3. For each $n \in D$, search for a monic polynomial P_n of degree n with

$$\partial_x^2 P_n + 2\omega\partial_x P_n + (\partial_x\omega + \omega^2 - r)P_n = 0. \tag{32}$$

If success is achieved then $\zeta_1 = P_n e^{\int \omega}$ is a solution of the differential equation. Else, case 1 cannot hold.

Case 2. Search for each $c \in \Gamma'$ and for ∞ the corresponding situation as follows:

Step 1. Search for each $c \in \Gamma'$ and ∞ the sets $E_c \neq \emptyset$ and $E_\infty \neq \emptyset$. For each $c \in \Gamma'$ and for ∞ we define $E_c \subset \mathbb{Z}$ and $E_\infty \subset \mathbb{Z}$ as follows:

(c_1): If $\circ\left(r_c\right) = 1$, then $E_c = \{4\}$
(c_2): If $\circ\left(r_c\right) = 2$, and $r = \cdots + b(x-c)^{-2} + \cdots,$ then

$$E_c = \left\{ 2 + k\sqrt{1+4b} : k = 0, \pm 2 \right\} \cap \mathbb{Z}.$$

(c_3): If $\circ\left(r_c\right) = v > 2$, then $E_c = \{v\}$
(∞_1): If $\circ\left(r_\infty\right) > 2$, then $E_\infty = \{0, 2, 4\}$
(∞_2): If $\circ\left(r_\infty\right) = 2$, and $r = \cdots + bx^2 + \cdots$, then

$$E_\infty = \left\{ 2 + k\sqrt{1+4b} : k = 0, \pm 2 \right\} \cap \mathbb{Z}.$$

(∞_3): If $\circ\left(r_\infty\right) = v < 2$, then $E_\infty = \{v\}$

Step 2. Find $D \neq \emptyset$ defined by

$$D = \left\{ n \in \mathbb{Z}_+ : \quad n = \frac{1}{2}\left(e_\infty - \sum_{c\in\Gamma'} e_c \right), \forall e_p \in E_p, \quad p \in \Gamma \right\}.$$

If $D = \emptyset$, then we should start the case 3. Now, if $\mathrm{Card}(D) > 0$, then for each $n \in D$ we search a rational function θ defined by

$$\theta = \frac{1}{2}\sum_{c\in\Gamma'} \frac{e_c}{x-c}.$$

Step 3. For each $n \in D$, search a monic polynomial P_n of degree n, such that

$$\partial_x^3 P_n + 3\theta\partial_x^2 P_n + (3\partial_x\theta + 3\theta^2 - 4r)\partial_x P_n + \left(\partial_x^2\theta + 3\theta\partial_x\theta + \theta^3 - 4r\theta - 2\partial_x r\right) P_n = 0. \tag{33}$$

If P_n does not exist, then case 2 cannot hold. If such a polynomial is found, set $\phi = \theta + \partial_x P_n / P_n$ and let ω be a solution of

$$\omega^2 + \phi\omega + \frac{1}{2}\left(\partial_x\phi + \phi^2 - 2r\right) = 0.$$

Then $\zeta_1 = e^{\int \omega}$ is a solution of the differential equation.

Case 3. Search for each $c \in \Gamma'$ and for ∞ the corresponding situation as follows:

Step 1. Search for each $c \in \Gamma'$ and ∞ the sets $E_c \neq \emptyset$ and $E_\infty \neq \emptyset$. For each $c \in \Gamma'$ and for ∞ we define $E_c \subset \mathbb{Z}$ and $E_\infty \subset \mathbb{Z}$ as follows:

(c_1): If $\circ(r_c) = 1$, then $E_c = \{12\}$
(c_2): If $\circ(r_c) = 2$, and $r = \cdots + b(x-c)^{-2} + \cdots$, then

$$E_c = \left\{6 + \frac{12k}{m}\sqrt{1+4b} : \quad k = 0, \pm 1, \ldots, \pm\frac{m}{2}, \quad m = 4, 6, 12\right\} \cap \mathbb{Z}.$$

(∞): If $\circ(r_\infty) = v \geq 2$, and $r = \cdots + bx^2 + \cdots$, then

$$E_\infty = \left\{6 + \frac{12k}{m}\sqrt{1+4b} : \quad k = 0, \pm 1, \ldots, \pm\frac{m}{2}, \quad m = 4, 6, 12\right\} \cap \mathbb{Z}.$$

Step 2. Find $D \neq \emptyset$ defined by

$$D = \left\{n \in \mathbb{Z}_+ : \quad n = \frac{m}{12}\left(e_\infty - \sum_{c \in \Gamma'} e_c\right), \forall e_p \in E_p, \quad p \in \Gamma\right\}.$$

In this case we start with $m = 4$ to obtain the solution, afterwards $m = 6$ and finally $m = 12$. If $D = \emptyset$, then the differential equation is not integrable because it falls in the case 4. Now, if $\mathrm{Card}(D) > 0$, then for each $n \in D$ with its respective m, search a rational function

$$\theta = \frac{m}{12}\sum_{c \in \Gamma'} \frac{e_c}{x-c}$$

and a polynomial S defined as

$$S = \prod_{c \in \Gamma'} (x-c).$$

Step 3. Search for each $n \in D$, with its respective m, a monic polynomial $P_n = P$ of degree n, such that its coefficients can be determined recursively by

$$P_{-1} = 0, \quad P_m = -P,$$

$$P_{i-1} = -S\partial_x P_i - \left((m-i)\,\partial_x S - S\theta\right) P_i - (m-i)\,(i+1)\,S^2 r P_{i+1},$$

where $i \in \{0, 1 \ldots, m-1, m\}$. If P does not exist, then the differential equation is not integrable because it falls in Case 4. Now, if P exists search ω such that

$$\sum_{i=0}^{m} \frac{S^i P}{(m-i)!}\omega^i = 0,$$

then a solution of the differential equation is given by

$$\zeta = e^{\int \omega},$$

where ω is solution of the previous polynomial of degree m.

A.2. Hamiltonian algebrization. In this section we follow [**1, 3**]. We recall that there are a lot of differential equations with coefficients that are not rational functions. For these differential equations it is useful, when is possible, to replace it by a new differential equation over the Riemann sphere $\mathbb{P}^1$ (that is, with rational coefficients). To do this, we can use a change of variables. The equation over $\mathbb{P}^1$ is called the *algebraic form* or *algebrization* of the original equation.

Hamiltonian change of variable. A change of variable $z = z(x)$ is called *Hamiltonian* if $(z(x), \partial_x z(x))$ is a solution curve of the autonomous classical one degree of freedom Hamiltonian system

$$\begin{array}{l}\partial_x z = \partial_w H\\ \partial_x w = -\partial_z H\end{array} \quad \text{with} \quad H = H(z,w) = \frac{w^2}{2} + V(z),$$

for some $V \in K$, where K is a differential field. Thus, $z = z(x)$ is a Hamiltonian change of variable if there exists $\alpha \in K$ such that $(\partial_x z)^2 = \alpha(z)$. More specifically, if $z = z(x)$ is a Hamiltonian change of variable, we can write $\partial_x z = \sqrt{\alpha}$, which leads us to the following notation: $\widehat{\partial}_z = \sqrt{\alpha}\partial_z$.

We can see that $\widehat{\partial}_z$ is a *derivation* because satisfy $\widehat{\partial}_z(f+g) = \widehat{\partial}_z f + \widehat{\partial}_z g$ and the Leibnitz rules

$$\widehat{\partial}_z(f\cdot g) = \widehat{\partial}_z f \cdot g + f \cdot \widehat{\partial}_z g, \quad \widehat{\partial}_z\left(\frac{f}{g}\right) = \frac{\widehat{\partial}_z f \cdot g - f \cdot \widehat{\partial}_z g}{g^2}.$$

We can notice that the chain rule is given by $\widehat{\partial}_z(f \circ g) = \partial_g f \circ g \widehat{\partial}_z(g) \neq \widehat{\partial}_g f \circ g \widehat{\partial}_z(g)$. The iteration of $\widehat{\partial}_z$ is given by

$$\widehat{\partial}_z^0 = 1, \quad \widehat{\partial}_z = \sqrt{\alpha}\partial_z, \quad \widehat{\partial}_z^n = \sqrt{\alpha}\partial_z \widehat{\partial}_z^{n-1} = \underbrace{\sqrt{\alpha}\partial_z\left(\ldots\left(\sqrt{\alpha}\partial_z\right)\right)}_{n \text{ times } \sqrt{\alpha}\partial_z}.$$

We call *Hamiltonian Algebrization* to the algebrization process obtained by a Hamiltonian change of variable.

Hamiltonian Algebrization Theorem [**1, 3**]**.** Consider the systems of linear differential equations $[A]$ and $[\widehat{A}]$ given respectively by

$$\partial_x \mathbf{Y} = -A\mathbf{Y}, \quad \widehat{\partial}_z \widehat{\mathbf{Y}} = -\widehat{A}\widehat{\mathbf{Y}}, \quad A = [a_{ij}], \quad \widehat{A} = [\widehat{a}_{ij}], \quad \mathbf{Y} = [y_{i1}], \quad \widehat{\mathbf{Y}} = [\widehat{y}_{i1}],$$

where $a_{ij} \in K = \mathbb{C}(z(x), \partial_x(z(x)))$, $\widehat{a}_{ij} \in \mathbb{C}(z) \subseteq \widehat{K} = \mathbb{C}(z, \sqrt{\alpha})$, $1 \leq i \leq n$, $1 \leq j \leq n$, $a_{ij}(x) = \widehat{a}_{ij}(z(x))$ and $y_{i1}(x) = y_{i1}(z(x))$. Suppose that L and $\widehat{L}$ are the Picard-Vessiot extensions of $[A]$ and $[\widehat{A}]$ respectively. If the transformation φ is given by

$$\varphi: \begin{array}{l} x \mapsto z \\ a_{ij} \mapsto \widehat{a}_{ij} \\ y_{i1}(x) \mapsto \widehat{y}_{i1}(z(x)) \\ \partial_x \mapsto \widehat{\partial}_z \end{array},$$

then the following statements hold:

1. $K \simeq \widehat{K}$, $\quad (K, \partial_x) \simeq (\widehat{K}, \widehat{\partial}_z)$.
2. $\mathrm{DGal}(L/K) \simeq \mathrm{DGal}(\widehat{L}/\widehat{K}) \subset \mathrm{DGal}(\widehat{L}/\mathbb{C}(z))$.
3. $(\mathrm{DGal}(L/K))^0 \simeq (\mathrm{DGal}(\widehat{L}/\mathbb{C}(z))^0$.

A natural example of Hamiltonian Algebrization, and for instance of the introduction of the new derivative $\widehat{\partial}_z$, is the case of second order linear differential equations. Consider $\partial_x^2 y + a\partial_x y + by = 0$, using φ we obtain $\widehat{\partial}_z^2 \widehat{y} + \widehat{a}\widehat{\partial}_z \widehat{y} + \widehat{b}\widehat{y} = 0$, which is equivalent to

$$\alpha \partial_z^2 \widehat{y} + \left(\frac{\partial_x \alpha}{2} + \sqrt{\alpha}\widehat{a} \right) \partial_z \widehat{y} + \widehat{b}\widehat{y} = 0, \tag{34}$$

where $y(x) = \widehat{y}(z(x))$, $\widehat{a}(z(x)) = a(x)$ and $\widehat{b}(z(x)) = b(x)$.

In general, for $y(x) = \widehat{y}(z(x))$, the equation $F(\partial_x^n y, \ldots, y, x) = 0$ with coefficients given by $a_{i_k}(x)$ is transformed in the equation $\widehat{F}(\widehat{\partial}_z^n \widehat{y}, \ldots, \widehat{y}, z) = 0$ with coefficients given by $\widehat{a}_{i_k}(z)$, where $a_{i_k}(x) = \widehat{a}_{i_k}(z(x))$. In particular, for $\sqrt{\alpha} \in \mathbb{C}(z)$ and $\widehat{a}_{i_k} \in \mathbb{C}(z)$, the equation $\widehat{F}(\widehat{\partial}_z^n \widehat{y}, \ldots, \widehat{y}, z) = 0$ is the Hamiltonian algebrization of $F(\partial_x^n y, \ldots, y, x) = 0$. Now, if each derivation ∂_x has order even, with $\alpha(z)$ and $\widehat{a}_{i_k}(z)$ being rational functions, then the equation $F(\partial_x^n y, \ldots, y, x) = 0$ admits an Hamiltonian Algebrization for $z = z(x)$. An example, that illustrate this, is given by the following linear differential equation:

$$\partial_x^{2n} y + a_{n-1}(x)\partial_x^{2n-2} y + \ldots + a_2(x)\partial_x^4 y + a_1(x)\partial_x^2 y + a_0(x)y = 0.$$

Hamiltonian Algebrization Algorithm Theorem [1, 3]. In general is very difficult to find a suitable Hamiltonian change of variable, for this reason Hamiltonian Algebrization is a method or procedure (not an algorithm!). For specific families of differential equations we can obtain algorithms to apply Hamiltonian Algebrization, for example *any differential equation*

$$F(\partial_x^n y, \partial_x^{n-1} y, \ldots, \partial_x y, y; e^{\lambda_1 x}, \ldots, e^{\lambda_k x}) = 0, \quad \mu = p_i \lambda_i, \quad p_i \in \mathbb{Z}, \quad 1 \leq i \leq k,$$

admits Hamiltonian Algebrization if and only if

$$\frac{\lambda_i}{\lambda_j} \in \mathbb{Q}^*, \quad 1 \leq i \leq k, \quad 1 \leq j \leq k$$

and by means of the Hamiltonian change of variable $z = e^{\mu x}$ we obtain the equation

$$\widehat{F}(\widehat{\partial}_z^n \widehat{y}, \widehat{\partial}_z^{n-1}\widehat{y}, \ldots, \widehat{\partial}_z \widehat{y}, y; z) = 0$$

with coefficients in $\mathbb{C}(z)$.

References

[1] P. B. Acosta-Humanez, *"Galoisian Approach to Supersymmetric Quantum Mechanics. The integrability analysis of the Schrodinger equation by means of differential Galois theory",* VDM Verlag, Dr Müller, Berlin, 2010.

[2] P. B. Acosta-Humanez, J. T. Lázaro-Ochoa, J. J. Morales-Ruiz & Ch. Pantazi, *On the integrability of polynomial fields in the plane by means of Picard-Vessiot theory*, Preprint 2012 arXiv:1012.4796

[3] P. B. Acosta-Humanez, J. J. Morales-Ruiz & J.-A. Weil, *Galoisian Approach to integrability of Schrödinger Equation*, Reports on Mathematical Physics, **67** (2011), no. 3, 305–374.

[4] A. Angelow & D. A. Trifonov, *Schrödinger covariance states in anisotropic waveguides*, preprint ICTP, IC/95/44 Trieste, Italy, 1995.

[5] A. Angelow, *Light propagation in nonlinear waveguide and classical two-dimensional oscillator,* Physica A 256 (1998) 485–98

[6] Berry M V, *Classical adiabatic angles and quantum adiabatic phase,* J. Phys. A: Math. Gen. 18 (1985), 15–27

[7] R. Cordero-Soto, R. M. Lopez, E. Suazo & S. K. Suslov, *Propagator of a charged particle with a spin in uniform magnetic and perpendicular electric fields*, Lett. Math. Phys. **84** (2008) #2–3, 159–178.

[8] R. Cordero-Soto, E. Suazo & S. K. Suslov, *Models of damped oscillators in quantum mechanics*, Journal of Physical Mathematics **1** (2009), S090603 (16 pages).

[9] R. Cordero-Soto, E. Suazo & S. K. Suslov, *Quantum integrals of motion for variable quadratic Hamiltonians*, Annals of Physics **325** (2010) #10, 1884–1912; see also Preprint 2010 arXiv:0912.4900v9.

[10] R. Cordero-Soto & S. K. Suslov, *Time reversal for modified oscillators*, Theoretical and Mathematical Physics **162** (2010) #3, 286–316; see also Preprint 2009 arXiv:0808.3149v9.

[11] R. Cordero-Soto & S. K. Suslov, *The degenerate parametric oscillator and Ince's equation,* J. Phys. A: Math. Theor. **44** (2011) 015101

[12] V. V. Dodonov, I. A. Malkin & V. I. Man'ko, *Integrals of motion, Green functions, and coherent states of dynamical systems,* Int. J. Theor. Phys. 14 (1975) 37–54

[13] V. V. Dodonov & Man'ko, *Invariants and Evolution of Nonstationary Quantum Systems*. Lebedev Institute Proceedings vol 183 (1987) (Nauka: Moscow) pp 71–181 (in Russian)

[14] V. V. Dodonov & V. I. Man'ko, *Invariants and Evolution of Nonstationary Quantum Systems*. Lebedev Institute Proceedings vol 183 (1989) (Commack, NY: Nova Science) pp 103–261 (Engl. Transl.)

[15] R. P. Feynman & A. R. Hibbs, *Quantum Mechanics and Path Integrals* (New York: McGraw-Hill), 1965.

[16] D. Gómez Vergel & E. J. S. Villaseñor, *The time-dependent quantum harmonic oscillator revisited: applications to quantum theory* Ann. Phys. 324 (2009) 1360–85

[17] E. L. Ince, *Ordinary differential equations*, Dover, 1956.

[18] J. H. Hannay, *Angle variable holonomy in adiabatic excursion of an integrable Hamiltonian* J. Phys. A: Math. Gen 18 (1985) 221–30

[19] I. Kaplansky, *An introduction to differential algebra*, Hermann, 1957.

[20] E. Kolchin,*Differential Algebra and Algebraic Groups*, Academic Press, 1973.

[21] J. Kovacic, *An Algorithm for Solving Second Order Linear Homogeneous Differential Equations*, J. Symbolic Computation, **2** (1986), 3–43.

[22] Lanfear, Nathan & Suslov, Sergei K, *The time-dependent Schroedinger equation, Riccati equation and Airy functions*, Preprint 2009 arXiv:0903.3608

[23] Li J-F, Fang J-Y, Xiao F-L, Liu X-H and Wang C-Z, *Time evolution and squeezing of degenerate and non-degenerate coupled parametric down-conversion with driving term Commun.* Theor. Phys. (Beijing, China) 51 (2009) 514–518

[24] C. F. Lo, *Coherent-state propagator of the generalized time-dependent parametric oscillator* Europhys. Lett. 24 (1993) 319–323

[25] R. M. Lopez & S. K. Suslov, *The Cauchy problem for a forced harmonic oscillator*, Preprint 2007 arXiv:0707.1902v8

[26] Louisell W H, *Radiation and Noise in Quantum Electronics* (New York: McGraw-Hill), 1964.

[27] Louisell W H and Yariv A, *Quantum fluctuations and noise in parametric processes: I.* Phys. Rev. 124 (1961) 1646–1654.

[28] Magnus W and Winkler S, *Hill's Equation* (New York: Dover), 1966.

[29] Malkin I A, Man'ko V I and Trifonov D A, *Linear adiabatic invariants and coherent states* J. Math. Phys. 14 (1973) 576–582

[30] J. Martinet & J.P. Ramis,*Théorie de Galois differentielle et resommation*, Computer algebra and differential equations, 117–214, Comput. Math. Appl., Academic Press, London, 1990.

[31] M. Meiler, R. Cordero–Soto, and S. K. Suslov, *Solution of the Cauchy problem for a time-dependent Schröodinger equation,* J. Math. Phys. 49 072102

[32] J. J. Morales-Ruiz, *"Differential Galois Theory and Non-integrability of Hamiltonian Systems"*, Progress in Mathematics 179, Birkhäuser, 1999.

[33] J. J. Morales-Ruiz and J. P. Ramis, *Galoisian obstructions to integrability of hamiltonian systems I & II*, Methods Appl. Anal. 8 (2001), no. 1, 33–95, 97–111.

[34] A.F. Nikiforov & V.B. Uvarov, *Special Functions of Mathematical Physics*, Birkhäuser, Basel, 198

[35] M. Orszag, *Quantum Optics* (Heidelberg: Springer), 1994.

[36] J. Peřina, Z. Hradil & B. Jurčo, *Quantum Optics and Fundamentals of Physics* (Dordrecht: Kluwer), 1994.

[37] H. Prakash, N. Chandra & Vachaspati, *Conversion of random-phase light into phase-coherent light* Phys. Rev. A 9 (1974) 2167–9

[38] R. R. Puri, *Mathematical Methods of Quantum Optics* (Berlin: Springer), 2001.

[39] M .T. Raiford, *Statistical dynamics of quantum oscillators and parametric amplification in a single mode.* Phys. Rev. A 2 (1970) 1541–58

[40] M. T. Raiford, *Degenerate parametric amplification with time-dependent pump amplitude and phase* Phys.Rev. A 9 (1974) 2060–9

[41] Y. R. Shen, *The Principles of Nonlinear Optics* (New York: Wiley), 1984.

[42] M. Studzinski & M. Przybylska, *Darboux Points and Integrability Analysis of Hamiltonian Systems with Homogeneous Rational Potentials*, Preprint 2012 arXiv:1205.4395

[43] E. Suazo & S.K. Suslov, *Soliton-like Solutions for Nonlinear Schroedinger Equation with Variable Quadratic Hamiltonians*, to appear on Journal of Russian Laser Research, 2012.

[44] E. Suazo & S. K. Suslov, *Cauchy problem for Schrödinger equation with variable quadratic Hamiltonians*, in preparation.

[45] S. K. Suslov, *Dynamical invariants for variable quadratic Hamiltonians*, Physica Scripta **81** (2010) #5, 055006 (11 pp); see also Preprint 2010 arXiv:1002.0144v6

[46] S. K. Suslov, *On the integrability of nonautonomous nonlinear Schrödinger equations*, Proc. Amer. Math. Soc. 140 (2012) #9, 3067-3082.

[47] H. Takahasi, *Information theory of quantum-mechanical channels Advances in Communication Systems: Theory and Applications vol 1 ed* A V Balakrishnan (New York: Academic) pp 227–310

[48] M. Van der Put & M. Singer, *Galois Theory in Linear Differential Equations*, Springer Verlag, New York, 2003.

[49] D. F. Walls & G. J. Milburn, *Quantum Optics* (Berlin: Springer), 1994.

[50] K-H Yeon, K-K Lee, Ch. I. Um, T. F. George & L. N. Pandey 1993 *Exact quantum theory of a time-dependent bound Hamiltonian systems* Phys. Rev. A 48, (1993) 2716–20

(P. Acosta-Humánez) Departamento de Matemáticas y Estadística Universidad del Norte Km. 5 via Puerto Colombia, Barranquilla, Colombia
E-mail address, P. Acosta-Humánez: `pacostahumanez@uninorte.edu.co`

(E. Suazo) University of Puerto Rico at Mayagüez Department of Mathematical Sciences CALL BOX 9000 Mayagüez, Puerto Rico 00681-9018
E-mail address, E. Suazo: `erwin.suazo@upr.edu`